

Math 218D: Week 1 Discussion

STUDY COPY

September 1, 2022

Problem 1. Fill in the blanks below.

$$A = \begin{bmatrix} 3 & -4 & 2 & 8 & 0 \\ 1 & 8 & 1/9 & 0 & 1 \\ 2 & 4 & -1 & \pi & 2 \end{bmatrix} \quad a_{23} = \underline{\hspace{1cm}} \quad \text{Col}_2(A) \in \mathbb{R} \text{---} \quad A^T = \underline{\hspace{3cm}}$$

Problem 2. Fill in the blanks below, assuming that S is *symmetric*.

$$S = \begin{bmatrix} 5 & -4 & \underline{\hspace{1cm}} & \underline{\hspace{1cm}} \\ \underline{\hspace{1cm}} & 19 & \underline{\hspace{1cm}} & -1 \\ 11 & 2 & 8 & 3 \\ 9 & \underline{\hspace{1cm}} & \underline{\hspace{1cm}} & -10 \end{bmatrix} \quad \text{trace}(S) = \underline{\hspace{1cm}}$$

Problem 3. By definition, a matrix S is *symmetric* if $\underline{\hspace{2cm}}$.

Problem 4. Suppose that A is $n \times n$ and let $S = A + A^T$. Prove that S is symmetric.

Hint. This proof can be quickly accomplished by filling in the blanks below.

$$\underline{\hspace{2cm}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}} = \underline{\hspace{2cm}}$$

Problem 5. Consider the matrix R given by

$$R = \begin{bmatrix} 1 & -3 & 0 & -9 & 5 \\ 0 & 0 & 1 & 14 & 9 \\ 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{Col}_4(R) = \underline{\hspace{1cm}} \text{Col}_1(R) + \underline{\hspace{1cm}} \text{Col}_3(R)$$

- (a) Fill in the blanks above to express the fourth column of R as a linear combination of the first and third columns of R .
- (b) Can the fifth column of R be expressed as a linear combination of the first and third columns of R ? Explain why or why not.

Problem 6. We write $\mathbb{R}^9 \xrightarrow{A} \mathbb{R}^{22}$ to indicate that A is a $___ \times ___$ matrix.

Problem 7. Suppose $\mathbb{R}^{13} \xrightarrow{M^T} \mathbb{R}^{37}$. Then M is a $___ \times ___$ matrix.

Problem 8. Fill in the blanks in the two equations below.

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = ___ \begin{bmatrix} ___ \\ ___ \end{bmatrix} + ___ \begin{bmatrix} ___ \\ ___ \end{bmatrix} + ___ \begin{bmatrix} ___ \\ ___ \end{bmatrix} \quad \begin{bmatrix} ___ & ___ \\ ___ & ___ \\ ___ & ___ \end{bmatrix} \begin{bmatrix} ___ \\ ___ \end{bmatrix} = 11 \begin{bmatrix} -3 \\ 5 \\ 1 \end{bmatrix} - 42 \begin{bmatrix} 7 \\ 6 \\ 4 \end{bmatrix}$$

Problem 9. Fill in the blanks in each equation below.

$$\begin{bmatrix} ___ \\ A \end{bmatrix} \begin{bmatrix} ___ \\ ___ \\ ___ \end{bmatrix} = \text{the third column of } A \quad \begin{bmatrix} ___ \\ A \end{bmatrix} \begin{bmatrix} ___ \\ ___ \\ ___ \end{bmatrix} = \text{the first column minus the third column of } A$$

$$\begin{bmatrix} ___ \\ A \end{bmatrix} \begin{bmatrix} ___ \\ ___ \\ ___ \end{bmatrix} = \text{the sum of all columns of } A \quad \begin{bmatrix} ___ \\ A \end{bmatrix} \begin{bmatrix} ___ \\ ___ \\ ___ \end{bmatrix} = \text{twice the first column of } A$$

Problem 10. Suppose that A has four columns. Fill in the blanks in the equation on the right to validate the equation on the left.

$$\text{Col}_1 + 3 \text{ Col}_2 - 9 \text{ Col}_3 = 6 \text{ Col}_4 \quad \begin{bmatrix} ___ \\ A \end{bmatrix} \begin{bmatrix} ___ \\ ___ \\ ___ \\ ___ \end{bmatrix} = \mathbf{0}$$

Problem 11. Find the missing entries in $A = \begin{bmatrix} * & 2 & -1 \\ -12 & -4 & 2 \\ * & 10 & -5 \\ * & 8 & -4 \end{bmatrix}$ assuming A has rank one.

Problem 12. We say that \mathbf{v} is an *eigenvector* of A with *corresponding eigenvalue* λ if $A\mathbf{v} = ______$.

Problem 13. Suppose that A is $n \times n$ and that $\mathbf{v} \in \mathcal{E}_A(\lambda)$. Calculate $(\lambda \cdot I_n - A)\mathbf{v}$.

Math 218D: Week 2 Discussion

STUDY COPY

September 8, 2022

Problem 1. $\langle [1 \ -3 \ 0 \ 2]^\top, [2 \ 1 \ 5 \ 0]^\top \rangle = \underline{\hspace{2cm}}$

Problem 2. Which of the following vectors is *orthogonal* to $\mathbf{v} = [1 \ 1 \ 1 \ 1]^\top$?

☐ $\mathbf{w} = [3 \ -5 \ 2 \ 1 \ 0]^\top$ ☐ $\mathbf{x} = [9 \ 2 \ 3 \ -6 \ -8]^\top$ ☐ $\mathbf{y} = [-1 \ -1 \ 9 \ -10 \ 3]^\top$


Problem 3. The length of \mathbf{v} can be calculated with an inner product using the formula $\|\mathbf{v}\| = \underline{\hspace{2cm}}$.

Problem 4. The inner product can be interpreted geometrically with the formula $\langle \mathbf{v}, \mathbf{w} \rangle = \underline{\hspace{2cm}}$.

Problem 5. If we view $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ as $n \times 1$ matrices, then $\langle \mathbf{v}, \mathbf{w} \rangle$ can be calculated using matrix multiplication with the formula $\langle \mathbf{v}, \mathbf{w} \rangle = \underline{\hspace{2cm}}$.

Problem 6. The adjoint formula for inner products states that $\langle A\mathbf{v}, \mathbf{w} \rangle = \underline{\hspace{2cm}}$.

Problem 7. Suppose that A and B are matrices satisfying $A^\top B = I_n$ and that \mathbf{v} and \mathbf{w} vectors making the following diagrams accurate.



Calculate $\langle B\mathbf{v} - 3\mathbf{w}, 2A\mathbf{v} - A\mathbf{w} \rangle$.

Problem 8. One of the following calculations is possible and the other is not. Carry out the possible calculation.

$$\begin{bmatrix} 5 & -1 \\ 1 & 1 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -2 & 1 \\ 3 & -3 \end{bmatrix} = \underline{\hspace{2cm}} \qquad \begin{bmatrix} 5 & -1 \\ 1 & 1 \\ 3 & 0 \end{bmatrix} \begin{bmatrix} 0 & -2 & 3 \\ 1 & 1 & -3 \end{bmatrix} = \underline{\hspace{2cm}}$$

Problem 9. Fill in the blanks in each of the following two equations.

$$\begin{bmatrix} _ & _ & _ \\ _ & _ & _ \\ _ & _ & _ \end{bmatrix} \begin{bmatrix} 0 & -3 & 1 & 0 \\ 1 & 9 & 0 & 0 \\ 0 & 4 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 9 & 50 & 5 & -4 \\ 12 & 193 & -3 & 19 \end{bmatrix} \qquad \begin{bmatrix} _ & _ \\ _ & _ \\ _ & _ \\ _ & _ \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 9 \\ 2 & -5 \\ 0 & 9 \end{bmatrix}$$

Problem 10. Suppose that A and B are 2022×2022 . Prove that $S = B^T A + A^T B$ is symmetric.

Problem 11. The last column of a matrix A is $[0 \ 3 \ 4]^T$ and the Gramian of A is

$$G = \begin{bmatrix} 9 & _ & -6 & _ \\ 3 & 14 & 13 & _ \\ _ & _ & 29 & _ \\ 5 & 13 & 0 & _ \end{bmatrix} = \underline{\hspace{2cm}}$$

- Fill in the missing entries of G and fill in the formula used to calculate G .
- The number of rows of A is $_$ and the number of columns of A is $_$.
- Which (if any) of the columns of A is orthogonal to the third column of A ?

Math 218D: Week 3 Discussion

STUDY COPY


September 15, 2022

Problem 1. Consider the system of equations given by

$$\begin{array}{rrrrrrrcl} x_1 & + & 2x_2 & - & 4x_3 & + & 9x_4 & = & -2 \\ 5x_1 & + & 11x_2 & - & 13x_3 & + & 37x_4 & = & 7 \\ -3x_1 & - & 6x_2 & + & 12x_3 & - & 24x_4 & = & 0 \end{array}$$

Use the Gauß-Jordan algorithm to find the general solution to this system.

Problem 2. Suppose A is a matrix satisfying $\text{rref}(A) = \begin{bmatrix} 1 & -13 & 0 & 6 \\ 0 & 0 & 1 & 17 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$. Fill-in the blanks below.



Problem 3. Use the Gauß-Jordan algorithm to calculate $\text{rref}(A)$ where $A = \begin{bmatrix} 3 & -6 & 12 & 0 & -9 \\ -7 & 14 & -28 & -5 & 26 \\ 5 & -12 & 12 & 2 & -13 \\ 2 & -3 & 12 & -3 & -5 \end{bmatrix}$.

Math 218D: Week 4 Discussion

STUDY COPY

September 22, 2022

Problem 1. Use the Gauß-Jordan algorithm to calculate $EA = R$ where $A = \begin{bmatrix} 1 & -5 & 0 & 3 \\ 0 & 0 & 5 & -5 \\ -5 & 25 & -11 & -4 \end{bmatrix}$.

Problem 2. Consider the $EA = R$ factorization and the vector \mathbf{b} given by

$$\begin{bmatrix} -6 & 5 & 2 & -13 \\ 4 & -4 & -2 & 9 \\ -9 & 9 & 4 & -21 \\ 1 & -2 & -1 & 3 \end{bmatrix} \begin{bmatrix} -3 & 21 & 2 & -26 & -5 & 3 \\ 3 & -21 & 1 & 14 & 38 & -2 \\ -3 & 21 & -3 & -6 & -60 & 1 \\ 2 & -14 & -1 & 16 & 7 & -2 \end{bmatrix} = \begin{bmatrix} 1 & -7 & 0 & 6 & 9 & 0 \\ 0 & 0 & 1 & -4 & 11 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \mathbf{b} = \begin{bmatrix} -1 \\ 0 \\ 2 \\ 1 \end{bmatrix}$$

Determine if $A\mathbf{x} = \mathbf{b}$ is consistent *without doing any row operations*.

Problem 3. Calculate $PA = LU$ where $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & -1 \\ 0 & -1 & -2 \\ 3 & -1 & 1 \end{bmatrix}$.

Problem 4. Consider the $PA = LU$ factorization and the vector \mathbf{b} given by

$$\begin{matrix} P \\ \left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{array} \right] \end{matrix} \begin{matrix} A \\ \left[\begin{array}{ccc} 1 & -1 & -1 \\ 1 & 0 & 0 \\ -1 & 1 & 1 \\ 3 & 1 & 3 \end{array} \right] \end{matrix} = \begin{matrix} L \\ \left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 3 & 4 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{array} \right] \end{matrix} \begin{matrix} U \\ \left[\begin{array}{ccc} 1 & -1 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{array} \right] \end{matrix} \quad \mathbf{b} = \begin{bmatrix} 15 \\ 11 \\ -15 \\ 13 \end{bmatrix}$$

Solve $A\mathbf{x} = \mathbf{b}$ without doing any row reductions.

Math 218D: Week 5 Discussion

STUDY COPY

September 29, 2022

Problem 1. To verify if $\mathbf{v} \in \text{Null}(A)$ we must check the equation _____.

Problem 2. Show that $\mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$ is in the null space of $A = \begin{bmatrix} 2 & 1 & 4 \\ 1 & 1 & 3 \\ 2 & 1 & 4 \\ 0 & 1 & 2 \end{bmatrix}$.

Problem 3. Suppose A is 5×9 and that $\mathbf{v} \in \text{Null}(A)$. Then $\mathbf{v} \in \mathbb{R}$ _____.

Problem 4. Suppose that A has four columns related by the equation $\text{Col}_4 = 3 \text{Col}_1 + \text{Col}_2 - \text{Col}_3$. Find a nonzero vector $\mathbf{v} \in \text{Null}(A)$.

Problem 5. A scalar λ is an *eigenvalue* of A if _____.

Problem 6. The *eigenspace* of A corresponding to an eigenvalue λ is $\mathcal{E}_A(\lambda) =$ _____.

Problem 7. An *eigenvector* $\mathbf{v} \in \mathcal{E}_A(\lambda)$ satisfies the equation _____.

Problem 8. Show that $\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$ is an eigenvector of $A = \begin{bmatrix} 10 & -6 & 6 \\ 6 & -2 & 6 \\ -6 & 6 & -2 \end{bmatrix}$ and identify the eigenvalue.

Problem 9. Find all vectors in $\mathcal{E}_A(-3)$ where $A = \begin{bmatrix} -23 & 40 & -60 \\ -5 & 7 & -15 \\ 5 & -10 & 12 \end{bmatrix}$.

Problem 10. To verify if $\mathbf{v} \in \text{Col}(A)$ we must check the equation _____.

Problem 11. Determine if $\mathbf{v} = \begin{bmatrix} 6 \\ 12 \\ 2 \end{bmatrix}$ is in the column space of $A = \begin{bmatrix} 1 & -3 & 4 \\ 2 & -6 & 8 \\ -3 & 9 & -12 \end{bmatrix}$.

Problem 12. To verify if $\mathbf{v} \in \text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ we must check _____.

Problem 13. Determine if $\begin{bmatrix} 3 & 0 & 1 \end{bmatrix}^\top \in \text{Span}\{\begin{bmatrix} 1 & -3 & -3 \end{bmatrix}^\top, \begin{bmatrix} -1 & 4 & 3 \end{bmatrix}^\top\}$.

Math 218D: Week 6 Discussion

STUDY COPY

October 6, 2022

Problem 1. By definition, what does it mean to call a list of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ *linearly dependent*?

Problem 2. By definition, what does it mean to call a list of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ *linearly independent*?

Problem 3. Determine if $\begin{bmatrix} 1 & -3 & 1 \end{bmatrix}^\top, \begin{bmatrix} -4 & 13 & -3 \end{bmatrix}^\top, \begin{bmatrix} 5 & -17 & 3 \end{bmatrix}^\top$ is independent.

Problem 4. Suppose that $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \in \mathbb{R}^n$ and let A be an $m \times n$ matrix such that $\{A\mathbf{v}_1, A\mathbf{v}_2, A\mathbf{v}_3\}$ is linearly independent. Show that $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly independent.

Problem 5. The columns of a matrix A are independent if and only if _____.

Problem 6. Suppose that A is a 3×3 matrix satisfying the following three equations.

$$\begin{bmatrix} A \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \\ -1 \end{bmatrix} \quad \text{rref} \left[\begin{array}{c|c} A & \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \end{array} \right] = \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right] \quad \text{rref} \left[\begin{array}{c|c} A^T & \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \end{array} \right] = \left[\begin{array}{ccc|c} 1 & 0 & 1 & -11 \\ 0 & 1 & 1 & -7 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Note that $\text{rref}(A)$ and $\text{rref}(A^T)$ can be inferred from the second and third equations above.

- (a) The vector $\begin{bmatrix} 1 & 1 & 1 \end{bmatrix}^T$ belongs to exactly one of the four fundamental subspaces of A . Select this space.
☐ The null space. ☐ The row space. ☐ The column space. ☐ The left null space.
- (b) The vector $\begin{bmatrix} 1 & -2 & -1 \end{bmatrix}^T$ belongs to exactly one of the four fundamental subspaces of A . Select this space.
☐ The null space. ☐ The row space. ☐ The column space. ☐ The left null space.
- (c) Determine if $\begin{bmatrix} 1 & -1 & 1 \end{bmatrix}^T \in \text{Null}(A)$.

Problem 7. Suppose that A and B are $n \times n$ matrices and that $\mathbf{v} \in \mathbb{R}^n$ satisfies $\mathbf{v} \in \mathcal{E}_A(-2)$ and $\mathbf{v} \in \mathcal{E}_B(5)$. Show that \mathbf{v} is an eigenvector of $M = A^2 + AB - I_n$ and identify the corresponding eigenvalue.

Math 218D: Week 7 Discussion

STUDY COPY

October 13, 2022

Problem 1. Consider the calculations

$$\text{rref} \begin{bmatrix} 9 & 4 & 4 \\ -36 & -16 & -16 \\ 20 & 9 & 4 \\ -49 & -22 & -12 \end{bmatrix}^A = \begin{bmatrix} 1 & 0 & 20 \\ 0 & 1 & -44 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{rref} \begin{bmatrix} 9 & -36 & 20 & -49 \\ 4 & -16 & 9 & -22 \\ 4 & -16 & 4 & -12 \end{bmatrix}^{A^T} = \begin{bmatrix} 1 & -4 & 0 & -1 \\ 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

(a) Find the pivot solutions to $A\mathbf{v} = \mathbf{0}$. These vectors form a basis of _____.

(b) Find the pivot solutions to $A^T\mathbf{v} = \mathbf{0}$. These vectors form a basis of _____.


(c) Find the pivot columns of A . These vectors form a basis of _____.

(d) Find the nonzero rows of $\text{rref}(A)$. These vectors form a basis of _____.

(e) The pivot columns of A^T form a basis of _____.

(f) The nonzero rows of $\text{rref}(A^T)$ form a basis of _____.

(g) Fill in the blanks in the figure below.



Problem 2. Suppose $EA = R$ where

$$E = \begin{bmatrix} 1 & -3 & -1 & 17 \\ -3 & 10 & 5 & -56 \\ 5 & -19 & -12 & 105 \\ -1 & 7 & 7 & -36 \end{bmatrix} \quad R = \begin{bmatrix} 1 & -3 & 0 & -7 & 0 \\ 0 & 0 & 1 & 9 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

(a) Draw the picture of the four fundamental subspaces of A , including their dimensions

(b) Find a basis of $\text{Col}(A^\top)$.

(c) Find a basis of $\text{Null}(A^\top)$.

(d) Find a basis of $\text{Null}(A)$.

(e) Find a basis of $\text{Col}(A)$.

Math 218D: Week 8 Discussion

STUDY COPY

October 20, 2022

Problem 1. Suppose that A is a matrix satisfying

$$\text{Col}(A^T) = \text{Span}\left\{\begin{bmatrix} 1 & 1 & 0 & 1 \end{bmatrix}^T, \begin{bmatrix} 0 & 3 & 2 & 4 \end{bmatrix}^T\right\} \quad \text{Null}(A^T) = \text{Span}\left\{\begin{bmatrix} 2 & 1 & 1 \end{bmatrix}^T\right\}$$

(a) Draw the picture of the four fundamental subspaces of A , including their dimensions

(b) Determine if $\mathbf{v} = \begin{bmatrix} 1 & 0 & 2 & -1 \end{bmatrix}^T$ satisfies $A\mathbf{v} = \mathbf{0}$.

(c) Determine if $\mathbf{b} = \begin{bmatrix} 3 & 5 & 2 \end{bmatrix}^T$ makes the system $A\mathbf{x} = \mathbf{b}$ consistent.


(d) Explain why $\text{Null}(A) \neq \text{Span}\left\{\begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}^T, \begin{bmatrix} 2 & 1 & 0 & 0 \end{bmatrix}^T\right\}$.

Problem 2. The *least squares problem* associated to $A\mathbf{x} = \mathbf{b}$ is _____.

Problem 3. Suppose $\hat{\mathbf{x}}$ is a least squares approximate solution to $A\mathbf{x} = \mathbf{b}$. Then $A\hat{\mathbf{x}} =$ _____.

Problem 4. The *least squares error* is defined as _____.

Problem 5. The figure below depicts the result of using the technique of least squares to fit a curve of the form $f(x) = c_0 + c_1 \cos(\pi x/3) + c_2 \sin(\pi x/3)$ to four data points.



Find the values of c_0 , c_1 , and c_2 and calculate the error in using $f(x)$ to approximate this data.

Math 218D: Week 9 Discussion

STUDY COPY

October 27, 2022

Problem 1. Define the concept of an $A = QR$ factorization.

Problem 2. A matrix M has orthonormal columns if and only if $M^\top M =$ _____.

Problem 3. Given $A = QR$, projection onto $\text{Col}(A)$ is given by $P_{\text{Col}(A)} =$ _____.

Problem 4. Suppose $A = QR$ where A has full column rank. Then the least squares problem $A^\top A \hat{x} = A^\top b$ reduces to _____.

Problem 5. Suppose that A is $m \times n$ with orthonormal columns and that $v \in \mathbb{R}^n$.

(a) Show that $\|Av\| = \|v\|$.

(b) Show that $n \leq m$.

Problem 6. Calculate $\begin{vmatrix} 1 & -9 & 3 & -8 \\ 9 & -81 & 27 & -70 \\ -5 & 45 & -14 & 29 \\ -7 & 60 & -16 & 52 \end{vmatrix}$.

Problem 7. Consider the following matrix factorization

$$\begin{bmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}^P \begin{bmatrix} 0 & -1 & -1 & 1 & 6 \\ -1 & -4 & 1 & -1 & -4 \\ -2 & -10 & 0 & 2 & -1 \\ 8 & 3 & 1 & -2 & -3 \\ 3 & -1 & 0 & 1 & 1 \end{bmatrix}^A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -1/4 & 1 & 0 & 0 & 0 \\ -1/8 & 29/74 & 1 & 0 & 0 \\ 0 & 4/37 & -1 & 1 & 0 \\ 3/8 & 17/74 & -8/19 & -12/19 & 1 \end{bmatrix}^L \begin{bmatrix} 8 & 3 & 1 & -2 & -3 \\ 0 & -37/4 & 1/4 & 3/2 & -7/4 \\ 0 & 0 & 38/37 & -68/37 & -273/74 \\ 0 & 0 & 0 & -1 & 5/2 \\ 0 & 0 & 0 & 0 & 97/38 \end{bmatrix}^U$$

Calculate $\det(A)$.

Problem 8. For $n \times n$ matrices A and B , $\det(A^T) =$ _____ and $\det(AB) =$ _____.

Problem 9. If possible, find 3×3 matrices A and B satisfying $\det(A + B) \neq \det(A) + \det(B)$. If this is not possible, then explain why.

Problem 10. The (i, j) minor of A is $M_{ij} =$ _____ and the (i, j) cofactor is $C_{ij} =$ _____.

Math 218D: Week 10 Discussion

STUDY COPY

November 3, 2022

Problem 1. Suppose that $\det(A) = 35$ and that each (i, j) minor of A is the (i, j) entry of $M = \begin{bmatrix} -45 & -9 & -41 & 5 \\ -10 & 5 & 15 & 5 \\ 45 & 16 & 34 & -5 \\ -10 & -2 & -13 & 5 \end{bmatrix}$.

(a) Find the cofactor matrix C of A and the adjugate matrix $\text{adj}(A)$.

(b) Find three independent vectors orthogonal to the first column of A .

(c) Solve $A\mathbf{x} = \mathbf{b}$ for $\mathbf{b} = [0 \ 7 \ 0 \ 0]^\top$.

Problem 2. Suppose that λ is an eigenvalue of an $n \times n$ matrix A . Then $\det(\lambda \cdot I_n - A) = \underline{\hspace{2cm}}$.

Problem 3. The reciprocal of $z = 7 - 9i$ is $1/z = \underline{\hspace{2cm}} + \underline{\hspace{2cm}}i$.

Problem 4. Consider the vectors $\mathbf{v} = [1 + i \quad 5]^\top$ and $\mathbf{w} = [1 - 3i \quad 2 + i]^\top$ and the matrix $A = \begin{bmatrix} 2 & 1+i \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ -2i & 1 \end{bmatrix}$.

(a) Calculate $\|\mathbf{v}\|$.

(b) Calculate $\langle \mathbf{v}, \mathbf{w} \rangle$.

(c) Calculate $A^* \mathbf{v}$.

Problem 5. We call a matrix A *Hermitian* if $A = A^*$. We call A *unitary* if $A^{-1} = A^*$.

Problem 6. Suppose that H is Hermitian. Show that every diagonal entry of H is a real number.

Problem 7. Suppose that U is $n \times n$ unitary and that $\mathbf{v}, \mathbf{w} \in \mathbb{C}^n$. Show that $\langle U\mathbf{v}, U\mathbf{w} \rangle = \langle \mathbf{v}, \mathbf{w} \rangle$.

Math 218D: Week 11 Discussion

STUDY COPY

November 10, 2022

Problem 1. The polynomial

$$f(t) = t^4 - 2t^3 - t^2 + t - 14$$

has four distinct roots r_1, r_2, r_3 , and r_4 .

(a) $r_1 + r_2 + r_3 + r_4 =$ _____ and $r_1 r_2 r_3 r_4 =$ _____

(b) Calculate $(1 - r_1)(1 - r_2)(1 - r_3)(1 - r_4)$.

Problem 2. Let r_1 and r_2 be the roots of

$$f(t) = -9t^2 - 2t - 1$$

Calculate $r_1^2 + r_2^2$.

Hint. Consider $(r_1 + r_2)^2$.

Problem 3. Consider the equation

$$\begin{bmatrix} 18 & 10 & -5 & 0 \\ -10 & * & 8 & 5 \\ -6 & -28 & -14 & -14 \\ -11 & -30 & -11 & -10 \end{bmatrix}^A = \begin{bmatrix} 18 & 10 & -5 & 0 \\ -10 & * & 8 & 5 \\ -6 & -28 & -14 & -14 \\ -11 & -30 & -11 & -10 \end{bmatrix}^X \begin{bmatrix} -1 & * & -7 & -1 \\ 0 & 0 & * & * \\ -1 & 6 & -1 & -6 \\ * & * & 2 & -1 \end{bmatrix}^B \begin{bmatrix} 11 & 10 & 0 & 3 \\ * & -7 & * & -3 \\ -10 & * & * & -3 \\ 3 & 3 & * & * \end{bmatrix}^{X^{-1}}$$

where the entries marked * are unknown. Find the missing entry of A .

Problem 4. Suppose that A has eigenspaces given by

$$\mathcal{E}_A(7) = \text{Span}\{[1 \ 3 \ 0]^\top\} \quad \mathcal{E}_A(1) = \text{Span}\{[-2 \ -5 \ -5]^\top\} \quad \mathcal{E}_A(-1) = \text{Span}\{[-3 \ -7 \ -9]^\top\}$$

Calculate $A^{2021}\mathbf{v}$ for $\mathbf{v} = [0 \ -1 \ 3]^\top$.

Math 218D: Week 12 Discussion

STUDY COPY

November 17, 2022

Problem 1. Consider the factorization

$$\begin{bmatrix} -233 & 693 \\ -84 & 250 \end{bmatrix}^A = \begin{bmatrix} 3 & 11 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 \\ 0 & 19 \end{bmatrix} \begin{bmatrix} 3 & 11 \\ 1 & 4 \end{bmatrix}^{-1}$$

(a) $\det(A) =$ _____

(b) Find the solution $\mathbf{u}(t)$ to the initial value problem $d\mathbf{u}/dt = A\mathbf{u}$ with $\mathbf{u}(0) = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$.

(c) Let V be the vector space consisting of all vectors \mathbf{v} such that the solution $\mathbf{u}(t)$ to $d\mathbf{u}/dt = A\mathbf{u}$ with $\mathbf{u}(0) = \mathbf{v}$ satisfies $\lim_{t \rightarrow \infty} \mathbf{u}(t) = \mathbf{0}$. Find a basis of V .

Problem 2. What is a spectral factorization? Which matrices have spectral factorizations?

Problem 3. Suppose S is a real-symmetric matrix whose eigenspaces are given by

$$\mathcal{E}_S(-3) = \text{Span}\left\{\begin{bmatrix} 1 & -2 & 0 & 2 \end{bmatrix}^\top, \begin{bmatrix} -1 & -3 & -2 & 2 \end{bmatrix}^\top\right\} \quad \mathcal{E}_S(5) = \text{Span}\left\{\begin{bmatrix} 0 & 2 & -1 & 2 \end{bmatrix}^\top\right\} \quad \mathcal{E}_S(9) = ?$$

(a) Find a basis of $\mathcal{E}_S(9)$.

(b) Find a spectral factorization of S .

Math 218D: Week 14 Discussion

STUDY COPY

December 1, 2022

Problem 1. What is a singular value decomposition?

Problem 2. Let σ_1 , σ_2 , and σ_3 be the singular values of

$$A = \begin{bmatrix} 2 & 0 & 1 & 1 \\ 2 & -2 & -1 & 0 \\ -3 & -1 & 0 & -1 \end{bmatrix}$$

Calculate $\sigma_1^2 + \sigma_2^2 + \sigma_3^2$.

Problem 3. If possible, construct a 2×2 matrix A with characteristic polynomial $\chi_A(t) = (t - 1)^2$ and with exactly one singular value given by $\sigma_1 = 1$. If this is not possible, then explain why.

Problem 4. Find the singular values of $A = \begin{bmatrix} 1 & -1 & 1 & 2 \\ -1 & 1 & -1 & -2 \\ 2 & -2 & 2 & 4 \end{bmatrix}$.

Problem 5. Suppose A is 3×3 with exactly two singular values $\sigma_1 = \sqrt{19}$ and $\sigma_2 = 3$. If possible, calculate $\det(A)$. If not, then explain why.

Problem 6. Suppose that $A = U\Sigma V^*$ is a singular value decomposition and define $A^+ = V\Sigma^{-1}U^*$. Show that $A^+\mathbf{b}$ is a least squares approximate solution to $A\mathbf{x} = \mathbf{b}$.

Math 218D: Week 15 Discussion

STUDY COPY

December 8, 2022

Problem 1. Let $\mathbf{f} : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be the function defined by

$$\mathbf{f}(x, y) = \begin{bmatrix} x^2 + \sin(x) \\ x(y - 2) \\ -3xy + y^2 \end{bmatrix}$$

Use the local linearization of \mathbf{f} at the point $P = (0, 1)$ to approximate $\mathbf{f}(1/2, 1/2)$.

Problem 2. Calculate $S = LDL^T$ where $S = \begin{bmatrix} 2 & 12 & 10 \\ 12 & 67 & 105 \\ 10 & 105 & -351 \end{bmatrix}$.

Problem 3. Suppose that factoring $S = LDL^\top$ allows us to write the quadratic form $q(\mathbf{x}) = \langle \mathbf{x}, S\mathbf{x} \rangle$ as

$$q(\mathbf{x}) = 10(x_1 - 5x_2 + 2x_3)^2 - 11(x_2 - 6x_3)^2 - 5x_3^2$$

Find L and D and determine the definiteness of S .

Problem 4. Determine the definiteness of $S = \begin{bmatrix} 0 & 4 & -6 & 8 & 16 \\ 4 & -650 & 3 & 1941 & -1 \\ -6 & 3 & 8 & -144 & 16 \\ 8 & 1941 & -144 & 2 & 18 \\ 16 & -1 & 16 & 18 & 52 \end{bmatrix}$.

Problem 5. Find R the Cholesky factorization $S = R^\top R$ of $S = \begin{bmatrix} 9 & 15 & -6 \\ 15 & 29 & 8 \\ -6 & 8 & 206 \end{bmatrix}$.

Math 218D: Week 9 Discussion

STUDY COPY

October 27, 2022

Problem 1. Define the concept of an $A = QR$ factorization.

Problem 2. A matrix M has orthonormal columns if and only if $M^\top M =$ _____.

Problem 3. Given $A = QR$, projection onto $\text{Col}(A)$ is given by $P_{\text{Col}(A)} =$ _____.

Problem 4. Suppose $A = QR$ where A has full column rank. Then the least squares problem $A^\top A \hat{\mathbf{x}} = A^\top \mathbf{b}$ reduces to _____.

Problem 5. Suppose that A is $m \times n$ with orthonormal columns and that $\mathbf{v} \in \mathbb{R}^n$.

(a) Show that $\|A\mathbf{v}\| = \|\mathbf{v}\|$.

(b) Show that $n \leq m$.

Problem 6. Calculate $\begin{vmatrix} 1 & -9 & 3 & -8 \\ 9 & -81 & 27 & -70 \\ -5 & 45 & -14 & 29 \\ -7 & 60 & -16 & 52 \end{vmatrix}$.

Problem 7. Consider the following matrix factorization

$$\begin{bmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}^P \begin{bmatrix} 0 & -1 & -1 & 1 & 6 \\ -1 & -4 & 1 & -1 & -4 \\ -2 & -10 & 0 & 2 & -1 \\ 8 & 3 & 1 & -2 & -3 \\ 3 & -1 & 0 & 1 & 1 \end{bmatrix}^A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -1/4 & 1 & 0 & 0 & 0 \\ -1/8 & 29/74 & 1 & 0 & 0 \\ 0 & 4/37 & -1 & 1 & 0 \\ 3/8 & 17/74 & -8/19 & -12/19 & 1 \end{bmatrix}^L \begin{bmatrix} 8 & 3 & 1 & -2 & -3 \\ 0 & -37/4 & 1/4 & 3/2 & -7/4 \\ 0 & 0 & 38/37 & -68/37 & -273/74 \\ 0 & 0 & 0 & -1 & 5/2 \\ 0 & 0 & 0 & 0 & 97/38 \end{bmatrix}^U$$

Calculate $\det(A)$.

Problem 8. For $n \times n$ matrices A and B , $\det(A^\top) =$ _____ and $\det(AB) =$ _____.

Problem 9. If possible, find 3×3 matrices A and B satisfying $\det(A + B) \neq \det(A) + \det(B)$. If this is not possible, then explain why.

Problem 10. The (i, j) minor of A is $M_{ij} =$ _____ and the (i, j) cofactor is $C_{ij} =$ _____.

Math 218D: Week 10 Discussion

STUDY COPY

November 3, 2022

Problem 1. Suppose that $\det(A) = 35$ and that each (i, j) minor of A is the (i, j) entry of $M = \begin{bmatrix} -45 & -9 & -41 & 5 \\ -10 & 5 & 15 & 5 \\ 45 & 16 & 34 & -5 \\ -10 & -2 & -13 & 5 \end{bmatrix}$.

(a) Find the cofactor matrix C of A and the adjugate matrix $\text{adj}(A)$.

(b) Find three independent vectors orthogonal to the first column of A .

(c) Solve $A\mathbf{x} = \mathbf{b}$ for $\mathbf{b} = [0 \ 7 \ 0 \ 0]^\top$.

Problem 2. Suppose that λ is an eigenvalue of an $n \times n$ matrix A . Then $\det(\lambda \cdot I_n - A) = \underline{\hspace{2cm}}$.

Problem 3. The reciprocal of $z = 7 - 9i$ is $1/z = \underline{\hspace{2cm}} + \underline{\hspace{2cm}}i$.

Problem 4. Consider the vectors $\mathbf{v} = [1 + i \quad 5]^\top$ and $\mathbf{w} = [1 - 3i \quad 2 + i]^\top$ and the matrix $A = \begin{bmatrix} 2 & 1+i \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ -2i & 1 \end{bmatrix}$.

(a) Calculate $\|\mathbf{v}\|$.

(b) Calculate $\langle \mathbf{v}, \mathbf{w} \rangle$.

(c) Calculate $A^* \mathbf{v}$.

Problem 5. We call a matrix A *Hermitian* if $A = A^*$. We call A *unitary* if $A^{-1} = A^*$.

Problem 6. Suppose that H is Hermitian. Show that every diagonal entry of H is a real number.

Problem 7. Suppose that U is $n \times n$ unitary and that $\mathbf{v}, \mathbf{w} \in \mathbb{C}^n$. Show that $\langle U\mathbf{v}, U\mathbf{w} \rangle = \langle \mathbf{v}, \mathbf{w} \rangle$.

Math 218D: Week 11 Discussion

STUDY COPY

November 10, 2022

Problem 1. The polynomial

$$f(t) = t^4 - 2t^3 - t^2 + t - 14$$

has four distinct roots r_1, r_2, r_3 , and r_4 .

(a) $r_1 + r_2 + r_3 + r_4 =$ _____ and $r_1 r_2 r_3 r_4 =$ _____

(b) Calculate $(1 - r_1)(1 - r_2)(1 - r_3)(1 - r_4)$.

Problem 2. Let r_1 and r_2 be the roots of

$$f(t) = -9t^2 - 2t - 1$$

Calculate $r_1^2 + r_2^2$.

Hint. Consider $(r_1 + r_2)^2$.

Problem 3. Consider the equation

$$\begin{bmatrix} 18 & 10 & -5 & 0 \\ -10 & * & 8 & 5 \\ -6 & -28 & -14 & -14 \\ -11 & -30 & -11 & -10 \end{bmatrix}^A = \begin{bmatrix} 18 & 10 & -5 & 0 \\ -10 & * & 8 & 5 \\ -6 & -28 & -14 & -14 \\ -11 & -30 & -11 & -10 \end{bmatrix}^X \begin{bmatrix} -1 & * & -7 & -1 \\ 0 & 0 & * & * \\ -1 & 6 & -1 & -6 \\ * & * & 2 & -1 \end{bmatrix}^B \begin{bmatrix} 11 & 10 & 0 & 3 \\ * & -7 & * & -3 \\ -10 & * & * & -3 \\ 3 & 3 & * & * \end{bmatrix}^{X^{-1}}$$

where the entries marked * are unknown. Find the missing entry of A .

Problem 4. Suppose that A has eigenspaces given by

$$\mathcal{E}_A(7) = \text{Span}\{[1 \ 3 \ 0]^\top\} \quad \mathcal{E}_A(1) = \text{Span}\{[-2 \ -5 \ -5]^\top\} \quad \mathcal{E}_A(-1) = \text{Span}\{[-3 \ -7 \ -9]^\top\}$$

Calculate $A^{2021}\mathbf{v}$ for $\mathbf{v} = [0 \ -1 \ 3]^\top$.

Math 218D: Week 12 Discussion

STUDY COPY

November 17, 2022

Problem 1. Consider the factorization

$$\begin{bmatrix} -233 & 693 \\ -84 & 250 \end{bmatrix}^A = \begin{bmatrix} 3 & 11 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 \\ 0 & 19 \end{bmatrix} \begin{bmatrix} 3 & 11 \\ 1 & 4 \end{bmatrix}^{-1}$$

(a) $\det(A) =$ _____

(b) Find the solution $\mathbf{u}(t)$ to the initial value problem $d\mathbf{u}/dt = A\mathbf{u}$ with $\mathbf{u}(0) = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$.

(c) Let V be the vector space consisting of all vectors \mathbf{v} such that the solution $\mathbf{u}(t)$ to $d\mathbf{u}/dt = A\mathbf{u}$ with $\mathbf{u}(0) = \mathbf{v}$ satisfies $\lim_{t \rightarrow \infty} \mathbf{u}(t) = \mathbf{0}$. Find a basis of V .

Problem 2. What is a spectral factorization? Which matrices have spectral factorizations?

Problem 3. Suppose S is a real-symmetric matrix whose eigenspaces are given by

$$\mathcal{E}_S(-3) = \text{Span}\left\{\begin{bmatrix} 1 & -2 & 0 & 2 \end{bmatrix}^\top, \begin{bmatrix} -1 & -3 & -2 & 2 \end{bmatrix}^\top\right\} \quad \mathcal{E}_S(5) = \text{Span}\left\{\begin{bmatrix} 0 & 2 & -1 & 2 \end{bmatrix}^\top\right\} \quad \mathcal{E}_S(9) = ?$$

(a) Find a basis of $\mathcal{E}_S(9)$.

(b) Find a spectral factorization of S .